



(12)

(21) 2 307 960

(22) 04.05.2000

(51) Int. Cl.<sup>7</sup>:

C12N 15/54, A01H 5/00,  
C12P 21/00, C12N 5/04,  
A01H 5/10, C12N 5/10,  
C12N 15/82

(30) 60/132,849 US 06.05.1999

(71)

**BOARD OF TRUSTEES OPERATING MICHIGAN  
STATE UNIVERSITY,  
238 Administration Building, EAST  
LANSING, XX (US).**

(72)

**BENNING, CHRISTOPH (US).**  
**DOERMAN, PETER (US).**

(74)


**DIMOCK STRATTON CLARIZIO LLP**

(54) ADN CODANT POUR LA DIGALACTOSYLDIACYLGLYCEROL GALACTOSYLTRANSFERASE VEGETALE ET METHODES D'UTILISATION

(54) METHODS FOR GENERATION  
DNA ENCODING FOR PLANT DIGALACTOSYLDIACYLGLYCEROL GALACTOSYLTRANSFERASE AND  
METHODS OF USE

(57)


The cDNA encoding digalactosyldiacylglycerol galactosyltransferase (DGD1) is provided. The deduced amino acid sequence is also provided. Methods of making and using DGD1 to screen for new herbicides and alter a plant's leaf lipid composition are also provided, as well as expression vectors, transgenic plants or other organisms transfected with said vectors.





(21) (A1) **2,307,960**  
(22) 2000/05/04  
(43) 2000/11/06

- (72) BENNING, CHRISTOPH, US  
(72) DOERMAN, PETER, US  
(71) BOARD OF TRUSTEES OPERATING MICHIGAN STATE  
UNIVERSITY, US  
(51) Int. Cl.<sup>7</sup> C12N 15/54, C12N 15/82, C12N 5/10, A01H 5/10, C12N 5/04,  
C12P 21/00, A01H 5/00  
(30) 1999/05/06 (60/132,849) US  
(54) **ADN CODANT POUR LA DIGALACTOSYLDIACYLGLYCEROL  
GALACTOSYLTRANSFERASE VEGETALE ET METHODES  
D'UTILISATION**  
(54) **DNA ENCODING FOR PLANT  
DIGALACTOSYLDIACYLGLYCEROL  
GALACTOSYLTRANSFERASE AND METHODS OF USE**



(57) The cDNA encoding digalactosyldiacylglycerol galactosyltransferase (DGD1) is provided. The deduced amino acid sequence is also provided. Methods of making and using DGD1 to screen for new herbicides and alter a plant's leaf lipid composition are also provided, as well as expression vectors, transgenic plants or other organisms transfected with said vectors.



# **ABSTRACT**

The cDNA encoding digalactosyldiacylglycerol galactosyltransferase (DGD1) is provided. The deduced amino acid sequence is also provided. Methods of making and using DGD1 to screen for new herbicides and alter a plant's leaf lipid composition are also provided, as well as expression vectors, transgenic plants or other organisms transfected with said vectors.

5

**DNA ENCODING FOR PLANT DIGALACTOSYLDIACYLGLYCEROL  
GALACTOSYLTRANSFERASE AND METHODS OF USE**

5

**FIELD OF THE INVENTION**

The invention relates generally to plant galactolipids and more particularly  
10 the gene encoding digalactosyldiacylglycerol galactosyltransferase.

**BACKGROUND OF THE INVENTION**

The process of photosynthesis is the basis for all life on earth because it  
provides oxygen and ultimately converts inorganic matter into organic matter. The  
photosynthetic apparatus in plant cells is associated with a particular membrane  
15 system inside chloroplasts, the thylakoids. Four lipids are found to be associated  
with thylakoid membranes in plants and photosynthetic bacteria. Only one of  
them is a phospholipid, the ubiquitous phosphatidylglycerol. The other three are  
non-phosphorous diacylglycerol glycolipids with one or two galactose moieties or  
a sulfonic acid derivative of glucose attached to diacylglycerol. Browse, J. et al.,  
20 *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42:467 (1991); Joyard, J. et al., *Plant*  
*Physiol.* 118:715 (1998). The galactolipids constitute the bulk (close to 80%) of  
the thylakoid lipid matrix and within green plant parts, 70-80% of the lipids are  
associated with photosynthetic membranes. Taking into account that plants  
represent the major portion of the global bioorganic matter, it comes as no  
25 surprise that the two galactolipids, mono- and digalactosyldiacylglycerol, are the  
most abundant lipids in the biosphere. Most vegetables and fruits in human or  
animal diets are rich in galactolipids. Their breakdown products represent an  
important dietary source of galactose and polyunsaturated fatty acids. Ohlsson,  
L. et al., *J. Nutrition* 128:239 (1998); Andersson, L. et al., *J. Lipid Res.* 36:1392  
30 (1995). The elucidation of the pathway for galactolipid biosynthesis has been  
extremely challenging. Thylakoid membrane lipid biosynthesis in plants is highly  
complex bringing together carbohydrate and fatty acid metabolisms. There is a  
mesmerising number of molecular species for each thylakoid lipid due to the large  
number of combinatorial possibilities for fatty acid substituents. Even more  
35 dazzling, the biosynthesis of thylakoid lipids is not restricted to enzymes

associated with the chloroplast where galactolipids are found, but also involves enzymes in the endoplasmic reticulum (ER) (Figure 1). The mechanism for subcellular trafficking of lipid moieties from the ER that ultimately become incorporated into the thylakoid lipids inside the plastids poses one of the most

5 challenging enigmas of modern plant biochemistry. Molecular species of galactolipids containing diacylglycerol moieties derived from the plastid or the ER pathway can be distinguished based on their fatty acid composition. Heinz, E. et al., *Plant Physiol.* 72:273 (1983). Lipid moieties assembled inside the plastid carry preferentially a 16-carbon fatty acid in the *sn*2-position of diacylglycerol,

10 while lipids derived from the ER pathway contain an 18-carbon fatty acid in this position. This is due to different substrate specificities of the respective acyltransferases in the plastid and the ER. An extensive screening of different plant species revealed that the plastid pathway is dispensable in many plants. Mongrand, S. et al., *Phytochemistry* 49:1049 (1998). However, no naturally

15 occurring plant has been found, in which the ER pathway was non-functional. A mutant of *Arabidopsis*, *act1*, is partially blocked in the plastid pathway. Kunst, L. et al., *PNAS (USA)* 85:4143 (1988). This mutant is deficient in the acyltransferase which catalyses the biosynthesis of lysophosphatidic acid inside the plastid (Figure 1). Other mutants of *Arabidopsis* have been described that

20 affect the fatty acid and, thus, the molecular species composition of thylakoid lipids. Browse, J. et al., in *Arabidopsis*, E.M. Meyerowitz and C.R. Somerville, Eds. (Cold Spring Harbor Laboratory Press, New York) pp. 881-912 (1994). Most of these are deficient in fatty acid desaturases. However, the only higher plant mutant known to be directly affected in galactolipid assembly is the *dgd1*

25 mutant of *Arabidopsis*. Dormann, P. et al., *Plant Cell* 7:1801 (1995). In this mutant the relative amount of the digalactosyl lipid is reduced to 10% of wild type. It has already proven to be very valuable in assessing the importance of the digalactosyl lipid for the assembly and function of the photosynthetic membranes. Growth, chloroplast ultra structure, the composition and relative

30 ratios of different pigment protein complexes, the light utilization by the photosynthetic apparatus, and the import of proteins into chloroplasts are affected in the *dgd1* mutant. Hartel, H. et al., *Plant Physiol.* 115:1175 (1997); Reifarth, F. et al., *Biochemistry* 36:11769 (1997); Hartel, H. et al., *Plant Physiol. Biochem.* 36:407 (1998); Chen, L.-J. et al., *Plant J.* 16:33 (1998). In addition to

35 the reduction in the amount of galactolipid, the *dgd1* mutant also shows a

peculiar alteration in the fatty acid composition of the monogalactosyl lipid with a characteristic increase in the amount of molecular species containing 18-carbon fatty acids. The accumulation of these molecular species of the monogalactosyl lipid is consistent with their presumed precursor function in the biosynthesis of the digalactosyl lipid. Based on labelling experiments with isolated chloroplasts (van Besouw, A. et al., *Biochim. Biophys. Acta* 529:44 (1978); Hemmskerk, J.W.M. et al., *Plant Physiol.* 93:1286 (1990)), it has been proposed that one galactose moiety is transferred from one monogalactosyl lipid onto a second to form the digalactosyl lipid (Figure 1). The released diacylglycerol moiety is made available for further thylakoid lipid assembly with the bulk appearing in monogalactosyl lipid. As can be assumed from the fatty acid composition of the digalactosyl lipid in the wild type (Browse, J. et al., *Biochem. J.* 235:25 (1986)), the responsible enzyme is specific for molecular species derived from the ER. Accordingly, approximately equal amounts of ER-derived molecular species are found in the digalactosyl and monogalactosyl lipids (Figure 1). Therefore, it is expected that the disruption of digalactosyl lipid biosynthesis in the *dgd1* mutant also disturbs the assembly of other thylakoid lipids, in particular the ER-derived monogalactosyl lipid.

It would thus be desirable to provide the wild-type *DGD1* gene encoding for digalactosyldiacylglycerol galactosyltransferase (DGD1). It would also be desirable to isolate and purify the gene product. It would be further desirable to provide *in vitro* and *in vivo* assays to screen for new herbicides that inhibit the *DGD1* gene product. Galactolipids are unique to plants and other photosynthetic organisms. Therefore, in contrast to most herbicides currently in use, herbicides that inhibit galactolipid biosynthesis will not be toxic to animals, humans or microbial organisms in the soil.

It would also be desirable to control the digalactosyldiacylglycerol levels in plants by controlling the expression of the gene encoding for the DGD1 protein. It would further be desirable to transform plants using the gene in order to alter their lipid composition. An alteration in lipid composition would provide plants with an increased resistance to environmental factors such as, but not limited to, temperature stress and/or pathogen infection. It would further provide an increase in the yield of crop plants such as leafy vegetables.

#### SUMMARY OF THE INVENTION

The present invention provides a novel purified and isolated nucleic acid

sequence encoding digalactosyldiacylglycerol galactosyltransferase (DGD1). The cDNA encoding DGD1 is set forth SEQ ID NO: 1. The deduced amino acid sequence of DGD1 is also provided and set forth in SEQ ID NO: 2. The protein has a predicted molecular weight of 91.8 kDa and has some sequence similarity in the C-terminal portion to bacterial and plant glycosyltransferases.

Methods for making and using the cDNA encoding DGD1 are also provided. For example, wild-type *DGD1* can be used to produce recombinant DGD1 in bacteria or yeast. Such recombinant protein can be used in either an *in vivo* or *in vitro* assay to screen compounds for new herbicides. Additionally, *DGD1* may be used to alter a plant's leaf lipid composition thus altering sensitivity to environmental factors such as, but not limited to, temperature stress and/or pathogen infection and, in some cases, increase the yield of crop plants. Expression vectors containing the cDNA, transgenic plants and other organisms, *e.g.*, *E. coli*, transfected with said vectors, as well as seeds from said plants, are also provided by the present invention.

Additional aspects, advantages, and features of the present invention will become apparent from the following description, taken in conjunction with the accompanying drawings.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The various advantages of the present invention will become apparent to one skilled in the art by reading the following specification and by referencing the following drawings in which:

Figure 1 is a schematic illustrating galactolipid biosynthesis in *Arabidopsis*;

Figure 2 is a photograph of four-week old *Arabidopsis* plants showing the appearance of the wild type, *act1*, *dgd1* and *act1*, *dgd1* double mutants;

Figures 3A-3D are schematics illustrating the map-based cloning of the *DGD1* gene;

Figure 4 is a photograph of a thin-layer chromatograph showing reconstitution of the plant galactolipid biosynthetic pathway in *E. coli*.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The cDNA sequence encoding digalactosyldiacylglycerol galactosyltransferase (DGD1) is set forth in SEQ ID NO: 1. The deduced amino acid sequence is provided in SEQ ID NO: 2. The protein has a predicted mass of 91.8 kDa. Sequence comparisons show some similarity in the C-terminal portion to bacterial and plant glycosyltransferases.

A method for producing DGD1 in a host cell is also provided in the present invention. The method includes the steps of introducing an expression vector comprising a cDNA encoding DGD1 or a functional mutant thereof into a host cell and expressing the cDNA in an amount sufficient to permit purification of the DGD1. A vector may include a promoter that is functional in either a eukaryotic or prokaryotic cell. Preferably, the vector is introduced into a prokaryotic cell, such as *E. coli* that is routinely used for production of recombinantly produced proteins. Alternatively, the vector is introduced into a eukaryotic cell, such as *Saccharomyes cerevisiae* (yeast), that is also routinely used for the production of recombinantly produced proteins. It is further contemplated that DGD1 may be manufactured using standard synthetic methods.

The availability of large amounts of recombinant protein will permit the rapid screening of compounds to identify new herbicides. It will be appreciated that either a cell lysate, partially purified or purified recombinant DGD1 can be used in both *in vitro* and *in vivo* screening assays. It will also be appreciated that purified protein from a plant such as *Arabidopsis* is also contemplated within the present invention, and can also be used to screen for new compounds. For example, galactosyltransferase activity assay is provided wherein the amount of digalactosyldiacylglycerol (DGD) produced in *E. coli* expressing recombinant DGD1 and monogalactosyldiacylglycerol synthase is determined by thin layer chromatography. Thus, in a preferred embodiment, the polar lipids are extracted from the *E. coli* cells with one volume of 1M KCl and 0.2M H<sub>3</sub>PO<sub>4</sub> and 2 volumes of methanol/chloroform (1:1, v/v). In another embodiment, the extracted polar lipids are separated by thin layer chromatography on ammonium sulfate-impregnated silica plates developed in acetone/toluene/water (90:30:8, v/v/v). DGD lipid is then visualized by staining with  $\alpha$ -naphthol. It will be appreciated that *E. coli* expressing recombinant DGD1 can be exposed to various compounds and the effect of such treatment on DGD production assessed.

Once a compound is identified as an inhibitor of DGD1, mutagenesis can be used to create DGD1 mutants that show decreased or no sensitivity to the inhibitory compound. DGD1 mutants can be made by known methods such as, but not limited to, site-directed mutagenesis or random mutagenesis, followed by screening for an active DGD1 mutant. It will be appreciated that such a mutant gene would be suitable for overexpression in crop plants, conferring resistance to the selected inhibitor compound.

Furthermore, sequences of the present invention may be used to alter a plant's leaf lipid composition. Naturally-selected mutants of *Arabidopsis* with either decreased or increased expression of DGD1 show altered lipid and fatty acid composition. Altering a plant's leaf lipid composition can increase a plant's resistance to environmental factors such as, but not limited to, heat and/or cold stress, increase resistance to pathogen infection, and/or increase crop yield, especially of leafy vegetables such as lettuce. The method of altering leaf lipid composition of a plant includes the steps of introducing an expression vector comprising a cDNA encoding DGD1 or a functional mutant thereof, operably linked to a promoter functional in a plant cell into the cells of plant tissue and expressing the encoded protein in an amount effective to alter the leaf lipid composition. The level of expression can be increased by either combining the cDNA with a promoter that provides for a high level of expression, or by introducing multiple copies into the cell so that multiple copies are integrated into the genome of transformed plant cells. Once transformed cells exhibiting increased DGD1 activity are obtained, transgenic plants and seeds can then be regenerated therefrom, and evaluated for the stability of the inheritance of altered leaf lipid composition.

The *DGD1* nucleotide sequence may thus be fused to a gene or fragment thereof, which allows it to be expressed in a plant cell. The *DGD1* nucleotide sequence in combination with the gene or gene fragment, is referred to as an "expression vector" herein. It will be appreciated that the expression vectors of the present invention may contain any regulatory elements necessary and known to those skilled in the art for expression of DGD1. For example, such vectors may contain, but are not limited to, sequences such as promoters, operators and regulators, which are necessary for, and/or may enhance, the expression of DGD1.

The invention also provides the nucleic acid sequence from a shorter gene on chromosome 4 of *Arabidopsis thaliana* that has a high sequence similarity to *DGD1* (blast P score of 360). The nucleic acid sequence, *DGD2*, is set forth in SEQ ID NO: 3 (GenBank Accession No. AF058919). The deduced amino acid sequence DGD2 is provided in SEQ ID NO: 4. The gene, *DGD2* encodes a protein missing approximately 340 amino acids of the N-terminal portion as compared to DGD1, but shows similarity to the glycosyltransferase-like sequence part of DGD1. The amino acid sequence homology between DGD1 and DGD2 is 64.4% over 365 amino acids. This high degree of homology indicates that DGD2 would have the same activity and a similar function as DGD1 in plants. Therefore, it

will be appreciated that the *DGD2* gene can be used with the methods of the present invention. The predicted DGD1 protein contains an N-terminal transit peptide typical for proteins imported into the plastid. Furthermore, two strongly hydrophobic domains (amino acids 347-372 and 644-670) were found in the sequence. While not wishing to be bound by theory, this observation agrees with a proposed association of DGD1 with the plastid envelope membranes. Block, M.A. et al., *J. Biol. Chem.* 258:13281 (1983); Cline, K. et al., *Plant Physiol.* 71:366 (1983); Dorne, A.-J. et al., *FEBS Lett.* 145:30 (1982).

As referred to herein, the term "cDNA" is meant a nucleic acid, either naturally occurring or synthetic, which encodes a protein product. The term "nucleic acid" is intended to mean natural and/or synthetic linear, circular and sequential arrays of nucleotides and nucleosides, *e.g.*, cDNA, genomic DNA (gDNA), mRNA, and RNA, oligonucleotides, oligonucleosides, and derivatives thereof. The term "encoding" is intended to mean that the subject nucleic acid may be transcribed and translated into either the desired polypeptide or the subject protein in an appropriate expression system, *e.g.*, when the subject nucleic acid is linked to appropriate control sequences such as promoter and enhancer elements in a suitable vector (*e.g.*, an expression vector) and when the vector is introduced into an appropriate system or cell. As used herein, "polypeptide" refers to an amino acid sequence which comprises both full-length protein and fragments thereof.

As referred to herein, the term "capable of hybridizing under high stringency conditions" means annealing a strand of DNA complementary to the DNA of interest under highly stringent conditions. Likewise, "capable of hybridizing under low stringency conditions" refers to annealing a strand of DNA complementary to the DNA of interest under low stringency conditions. In the present invention, hybridizing under either high or low stringency conditions would involve hybridizing a nucleic acid sequence (*e.g.*, the complementary sequence to SEQ ID No. 1 or portion thereof), with a second target nucleic acid sequence. "High stringency conditions" for the annealing process may involve, for example, high temperature and/or lower salt content, which disfavor hydrogen bonding contacts among mismatched base pairs. "Low stringency conditions" would involve lower temperature, and/or higher salt concentration than that of high stringency conditions. Such conditions allow for two DNA strands to anneal if substantial, as is the case among DNA strands that code for the same protein but differ in

sequence due to the degeneracy of the genetic code. Appropriate stringency conditions which promote DNA hybridization, for example, 6X SSC at about 45° C, followed by a wash of 2X SSC at 50° C, are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1989), 6.31-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2X SSC at 50° C, to a high stringency of about 0.2X SSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency at room temperature, about 22° C, to high stringency conditions, at about 65° C. Other stringency parameters are described in Maniatis, T., et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring NY, (1982), at pp. 387-389; see also Sambrook J. et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Volume 2, Cold Spring Harbor Laboratory Press, Cold Spring, NY at pp. 8.46-8.47 (1989).

The foregoing and other aspects of the invention may be better understood in connection with the following examples, which are presented for purposes of illustration and not by way of limitation.

#### SPECIFIC EXAMPLE 1

##### CONSTRUCTION OF *act1, dgd1* DOUBLE MUTANT

An *act1, dgd1* double mutant (Figure 1) was constructed. An F<sub>2</sub> population of plants derived from the cross of *act1* (*Arabidopsis* Biological Resource Center, Columbus, Ohio) and *dgd1* was screened by thin-layer and gas chromatography for the characteristic lipid and fatty acid phenotype anticipated for the double mutant. About 1/16 of the F<sub>2</sub> plants contained strongly reduced amounts of 7,10,13-hexadecatrienoic acid as found in the *act1* mutant and reduced amounts of the galactolipid DGD, indicative for the *dgd1* mutation. This was the expected result for the segregation of two recessive unlinked mutations. Four-week-old representative plants raised on soil of the wild type (ecotype Col-2), the single homozygous mutants *act1* and *dgd1*, and the *act1, dgd1* double homozygous mutant are shown in Figure 2. The double mutant was severely stunted (Figure 2), and showed a more extreme growth phenotype than each of the mutant parents and any other known lipid mutant of *Arabidopsis*. The fatty acid composition of the galactolipids of the mutants as well as their fraction of total polar lipids were measured in leaves obtained from tissue-culture grown plants by thin-layer chromatography and subsequent gas chromatography of fatty acid

methyl esters (Table 1).

	Wild Type <sup>a</sup>	<i>dgd1</i> <sup>a</sup>	<i>act1</i> <sup>a</sup>	<i>act1,dgd1</i> <sup>a</sup>
Monogalactosyldiacylglycerol	49.8	53.5	53.7	44.2
7,10,13-hexadecatrienoic acid	28.9	14.1	0.8	0.7
$\alpha$ -linolenic acid	62.5	78.5	88.5	86.4
Digalactosyldiacylglycerol	12	1.7	15.9	2.1
7,10,13-hexadecatrienoic acid	2.6	2.8	n.d.	n.d.
$\alpha$ -linolenic acid	71.1	41.2	83.0	44.6

5       <sup>a</sup>The values are given in mol % of total polar leaf lipids for the two galactolipids and mol % of fatty acids attached to each of the two galactolipids for 7,10,13-hexadecatrienoic acid (*all-cis*-16:3 $\Delta^{7,10,13}$ ) and  $\alpha$ -linolenic acid (*all-cis*-18:3 $\Delta^{9,12,15}$ ). The values represent the means of three measurements each. The standard deviation was below 2.5% (galactolipids) and 1.0% (fatty acids). n.d., not detectable.

10       Because the lipid and fatty acid composition of the double mutant was not more severe than that of either of the parents, it is unlikely that the extreme growth phenotype of the double mutant is due to specific lipid or fatty acid effects. Thus, it seems plausible to conclude that the double mutant cannot produce sufficient amounts of thylakoid membranes, because the plastid pathway is affected by the *act1* mutation and the ER pathway by the *dgd1* mutation. A

15       pathway model consistent with the available biochemical data and the single and double mutant phenotypes is shown in Figure 1. For clarity, the model focuses on the galactolipids representing close to 80% of all thylakoid lipids. At least two genes encoding putative monogalactosyl lipid synthases are present in *Arabidopsis* (Shimojima, M. et al., *PNAS (USA)* 94:333 (1997); and it is

20       proposed that these have different substrate specificities and different association with the inner or outer envelope in accordance with previous studies. Block, M.A. et al., *J. Biol. Chem.* 258:13281 (1983); Cline, K. et al., *Plant Physiol.* 71:366 (1983). This enzyme class is currently under investigation by others. Shimojima, M. et al., *PNAS (USA)* 94:333 (1997); Teucher, T. et al.,

25       *Planta* 184:319 (1991); Marechal, E. et al., *Biol. Chem.* 269:5788 (1994). Accordingly, a transient pool of monogalactosyl lipid is produced at the outer envelope from ER-derived diacylglycerol and immediately converted by DGD1. This process is accompanied by a transfer of lipid moieties from the outer to the inner envelope. In the absence of DGD1, monogalactosyl lipid cannot be

30       efficiently synthesized via the ER-pathway but the plastid pathway can compensate for this deficiency. Only when both pathways are blocked as in the *act1,dgd1* double mutant, the overall galactolipid biosynthesis is reduced to

levels insufficient to support growth. Apparently, the proposed initial biosynthesis of galactolipids at the outer envelope membrane cannot compensate for the biosynthesis via DGD1, but would explain the small amount of digalactosyl lipid and the altered molecular species composition of monogalactosyl lipid in the *dgd1* mutant.

## SPECIFIC EXAMPLE 2

### ISOLATION AND PURIFICATION OF THE GENE ENCODING DGD1

While the analysis of the *dgd1* mutant and the *act1,dgd1* double mutant in Specific Example 1 revealed the crucial role of DGD1 in galactolipid biosynthesis and subcellular lipid trafficking in higher plants, only the molecular and biochemical analysis of the *dgd1* gene product will yield a true understanding of the underlying mechanism. The cloning of both the mutant *dgd1* and wild-type *DGD1* genes represents the first step in this direction. Because no molecular information was available, the *dgd1* locus and the corresponding *dgd1* cDNA was isolated by a strategy based on the map position of *dgd1*. One of the difficulties encountered was the heterogeneous genetic background in the *dgd1* mutant with markers characteristic for ecotypes Col-2 or *Ler* found interspersed around the *dgd1* locus. This problem was solved by integration of two different mapping populations derived from crosses of *dgd1* to Col-2 or *Ler* wild-type. From a total of 135 F<sub>2</sub> plants derived from the cross *dgd1* x Col-2, plants with cross-overs between the two PCR markers nga162 and nga172 were selected. Bell, C.J. et al., *Genomics* 19:137 (1994). In this F<sub>2</sub> population, the *dgd1* locus was mapped relative to the RFLP markers g4523, *fad7*, g4547, 5E-5 and 18A-1. Similarly, a total of 424 F<sub>2</sub> plants from the cross *dgd1* x *Ler* were screened for cross-overs between the PCR markers nga127 and ATHCHIB. This mapping population was used to score the RFLP markers g2488a and 31A-H. The RFLP markers were obtained from the *Arabidopsis* Biological Resource Center at Columbus, Ohio (g4523, *fad7*, g2488a, g4547) or from genomic fragments (31A-H, 5E-5, 18A-1) isolated from cosmid inserts in this study. Unambiguous scoring of the mutant phenotype had to be done by thin-layer chromatography of leaf lipid extracts, requiring several thousand samples to be processed during the fine mapping process. The map encompassing the *DGD1* locus and the YAC, BAC, and cosmid contigs spanning the locus on chromosome three are shown in Figure 3. Figure 3A shows the genetic map of the relevant part of *Arabidopsis* chromosome 3. Two YAC clones containing *dgd1* are shown. Numbers indicate

recombinations between a given marker and *dgd1* per number of chromosomes analyzed in the respective mapping population. Figure 3B shows the fine mapping between the markers *fad7* and g2488a. The BAC (IGF clone#) and cosmid (C clone#) contigs are shown. Complementing clones are marked by  $\oplus$ , non-complementing by  $-$ . Figure 3C is a map of the cosmids C49B, C5A and C5E with H indicating *Hind*III restriction sites. Finally, Figure 3D shows the structure of the *DGD1* gene and cDNA. The exons are shaded and numbered 1 to 7. The sequence predicted to be a chloroplast transit peptide (T) is indicated, as well as the part showing similarity to glycosyltransferases (GTF, cross hatched), the start and stop codon (ATG, TAG, respectively), and the C to T mutation observed in the *dgd1* mutant. Genomic DNA isolated from the markers *fad7*, g4547 and g2488a was used to isolate DNA fragments from different libraries (CIC Yeast Artificial Chromosome library (Camilleri, C. et al., *Plant J.* 14:633 (1998)); IGF Bacterial Artificial Chromosome library (Mozo, T. et al., *Mol. Gen. Genet.* 258:562 (1998)); *Arabidopsis* cosmid library (Meyer, K. et al., in *Genome Mapping in Plants*, A. H. Patterns, Ed. (Academic Press, New York, 1996), pp. 137-154). Different cosmids harbouring inserts between T-DNA borders were tested for complementation. Because the *dgd1* mutant could not be transformed with these large genomic fragments, the clones were transferred into the wild type first and crossed the T-DNA into the *dgd1* mutant. Cosmid clones were transferred into *Agrobacterium tumefaciens* C58C1(pGV2260) and used to transform *Arabidopsis thaliana* Col-2 wild type plants via vacuum infiltration (Bechtold, N. et al., *Acad. Sci. Paris Life Sci.* 316:1194 (1993); Bent, A.F. et al., *Science* 265:1856 (1994)). Transformants were crossed with *dgd1* mutant plants and the segregation pattern in the F<sub>2</sub> generation was analyzed. Complementation was assumed when of 100 tested F<sub>2</sub> plants carrying the T-DNA all were phenotypically wild-type, whereas in non-complementing lines, a segregation of the wild-type versus mutant phenotype of 3 : 1 was expected. A minimum of 100 transgenic F<sub>2</sub> plants derived from each cross (1 to 3 independent crosses per cosmid) were analyzed. To avoid the possibility that the construct was corrupted by chance in any particular plant, several independent transgenic lines were used. The analysis of three F<sub>2</sub> populations derived from crosses with independent lines containing cosmid C49B was consistent with genetic complementation by a gene encompassed by the insert. Several cosmids overlapping (C5A, C5E) or neighbouring C49B (C27A, C14B) did not complement

the mutation. Large portions of the cosmid C49B were sequenced (SEQ ID NO: 5). One putative gene was located in the centre of C49B, but was only partially contained by the cosmids C5A and C5E. Based on the complementation analysis for the cosmids C49B, C5A and C5E it was concluded that this gene represents the *DGD1* locus. Therefore, the insert of C49B was used to screen a cDNA library. Uwer, U. et al. *Plant Cell* 10:1277 (1998). A 2683 bp long cDNA was identified and sequenced (Figures 3D and SEQ ID NO: 1). This cDNA appeared to be complete because it contained in-frame stop codons 5'-prime of a putative ATG start codon. The cDNA was inserted behind a CaMV 35S promoter and transferred directly into the mutant by *Agrobacterium* mediated *in planta* transformation. For direct complementation analysis, the *DGD1* cDNA released from pBluescriptIIISK(+) with *Sma*I, *Xho*I was ligated into the *Sma*I, *Sa*I sites of the binary vector pBINAR-Hyg (Becker, D., *Nucl. Acids Res.* 18:203 (1990)) in sense orientation behind the CaMV 35S promoter. This construct was directly transferred into the *dgd1* mutant via *Agrobacterium* by vacuum infiltration. Two transgenic plants were recovered which were phenotypically wild type with regard to habitus and lipid composition indicating complementation. To obtain corroborating evidence for complementation and to exclude the possibility of wild-type contamination, genetically homozygous *dgd1* plants were identified in each complementation experiment by DNA/DNA hybridization using the RFLP marker 5E-5 which scores identical in the Col-2 and Ler wild type background but different in the *dgd1* mutant. These plants were tested for lipids. With no exception, cosmid C49B and the *DGD1* cDNA lead to wild-type lipid composition in all tested transgenic plants homozygous for *dgd1*. To obtain independent evidence for the identity of the *DGD1* locus, the respective genomic DNA of the wild-type *DGD1* and the mutant *dgd1* loci were sequenced (SEQ ID NOS: 5 and 6). Further comparison of the genomic and cDNA sequences revealed 7 exons and a transition of a CAA codon (glutamine) to TAA in exon 6 in the *dgd1* mutant gene leading to a premature stop codon.

30

### **SPECIFIC EXAMPLE 3**

#### **RECONSTITUTION OF THE PLANT GALACTOLIPID BIOSYNTHETIC PATHWAY IN *E. COLI***

The *DGD1* cDNA is predicted to encode a 91.8 kDa protein with some sequence similarity in the C-terminal portion to bacterial and plant glycosyltransferases. To determine the biosynthetic activity of the wild-type

35

*DGD1* gene product, the *DGD1* cDNA was expressed in *E. coli* along with the monogalactosyldiacylglycerol (MGD) synthase previously isolated from cucumber. Shimojima, M. et al., *PNAS (USA)* 94:333 (1997). A 459 bp *XhoI*, *PvuII* fragment including the expression cassette was isolated from pQE31  
5 (Qiagen Inc.) and ligated into the *Sall*, *EcoRV* sites of pACYC184 (Chang, A.C.Y. et al., *J. Bacteriol.* 134:1141 (1978)) giving rise to the plasmid pACYC-31. The open reading frame of the *DGD1* cDNA was amplified by PCR using the primers 5'-GCGGATCCGGTAAAGGAACTCTAATT-3' (Ben239; SEQ ID NO: 7) and 5'-TTCTGCAGTCTACCAGCCGAAGATTGG-3' (Ben241; SEQ ID NO: 8), thereby  
10 introducing a *BamHI* site at the 5' and a *PstI* site at the 3' terminus. This cDNA fragment was ligated into the corresponding restriction sites of pACYC-31. The resulting vector, pACYC-31/239, was transferred into XL1-Blue cells carrying the expression vector pGEX-3X with the cucumber MGD synthase cDNA (Shimojima et al. 1998). The cells were grown and protein expression induced with IPTG.  
15 The QIA expressionist: A handbook for high-level expression and purification of 6x His-tagged proteins. Qiagen, Inc., Valencia, CA (1997). The polar lipids were extracted from the *E. coli* cells with 1 volume 1 M KCl, 0.2 M H<sub>3</sub>PO<sub>4</sub> and 2 volumes methanol/chloroform (1:1, v/v). Polar lipids were then separated by thin-layer chromatography on ammonium sulfate (0.15 M) impragnated Baker  
20 Si250PA silica plates developed in acetone/toluene/water (90:30:8, v/v/v). Digalactosyldiacylglycerol was visualized by staining with  $\alpha$ -naphthol. In addition to MGD a new glycolipid was observed (Lane 2, Figure 4) that co-migrates with an authentic digalactosyldiacylglycerol (DGD) standard (Lane 3, Figure 4). In contrast, when the MGD synthase gene but not the *dgd1* cDNA was present, no  
25 DGD was observed (Lane 1, Figure 4). The plant galactolipid biosynthetic pathway was therefore, reconstituted in *E. coli*. Furthermore, this result demonstrates that the *dgd1* gene indeed encodes a DGD galactosyltransferase.

A gene essential for the biosynthesis of the thylakoid lipid digalactosyldiacylglycerol was isolated from *Arabidopsis* by map-based cloning.  
30 The *act1, dgd1* double mutant analysis strongly suggests that DGD1 also plays a critical role in lipid trafficking of ER-derived thylakoid lipids in higher plants. The availability of the wild-type *DGD1* gene, the similar *DGD2* gene, as well as genes encoding monogalactosyl lipid synthases of *Arabidopsis* will permit the rigorous testing of the current hypothesis for galactolipid biosynthesis and subcellular lipid  
35 trafficking described herein.

The foregoing discussion discloses and describes merely exemplary embodiments of the present invention. One skilled in the art will readily recognize from such discussion, and from the accompanying drawings, that various changes, modifications and variations can be made therein without departing from the spirit and scope of the invention.

5

Patent and literature references cited herein are incorporated by reference as if fully set forth.

SEQUENCE LISTING

<110> Benning, Christoph  
Doermann, Peter

<120> DNA Encoding For Plant Digalactosyldiacylglycerol  
Galactosyltransferase And Methods of Use

<130> 6550-000042

<140>

<141>

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 2649

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```
gaattcggca cgaggcagca ctctcacgaa atcgctcgtga cgaaacgata aaccctaagt 60
catggtaaag gaaactctaa ttctccgctc atctacgtca atgacgaccg gaacatcttc 120
ttcttcgtct ctttcaatga cgttatcctc aacaaacgcg ttatcgtttt tgtcgaaagg 180
atggagagag gtatgggatt cagcagatgc ggatttgcag ctgatgagag acagagctaa 240
ctctgttaag aatctagcat caacgttcga tagagagatc gagaatttcc tcaataactc 300
ggcgaggtct gcgtttcccg ttgggtcacc atcggcgctc tctttctcaa atgaaattgg 360
tatcatgaag aagcttcagc cgaagatttc ggagtttcgt aggggttatt cggcgccgga 420
gattagtcgc aaggttatgg agagatgggg acctgcgaga gcgaagcttg gaatggatct 480
atcggcgatt aagaaggcga ttgtgtctga gatggaattg gatgagcgctc agggagtttt 540
ggagatgagt agattgagga gacggcgtaa tagtgatagg gttaggttta cggagttttt 600
cgcgagggct gagagagatg gagaagctta tttcggtgat tgggaaccga ttaggtcttt 660
gaagagtaga tttaaagagt ttgagaaacg aagctcgtta gaaatattga gtggattcaa 720
gaacagtga tttgttgaga agctcaaaac cagctttaaa tcaatttaca aagaaactga 780
tgaggctaag gatgtccctc cgttgatgt acctgaactg ttggcatgtt tggttagaca 840
atctgaacct tttcttgatc agattggtgt tagaaaggat acatgtgacc gaatagtaga 900
aagcctttgc aaatgcaaga gccacaact ttggcgtctg ccatctgcac aagcatccga 960
tttaattgaa aatgataacc atggagtga tttggatatg aggatagcca gtgttcttca 1020
aagcacagga caccattatg atggtgggtt ttggactgat tttgtgaagc ctgagacacc 1080
ggaaaacaaa aggcattgtg caattgttac aacagctagt cttccttgga tgaccggaac 1140
agctgtaaat ccgctattca gagcggcgta tttggcaaaa gctgcaaaac agagtgttac 1200
tctcgtgggt ccttggtctc gcgaatctga tcaagaacta gtgtatccaa acaatctcac 1260
cttcagctca cctgaagaac aagagagtta tatacgtaaa tggttggagg aaaggattgg 1320
tttcaaggct gattttaaaa tctcctttta ccagggaaag ttttcaaaag aaaggcgag 1380
```

catatttcct gctggtgaca cttctcaatt tatatcgtca aaagatgctg acattgctat 1440  
 acttgaagaa cctgaacatc tcaactggta ttatcacggc aagcgttgga ctgataaatt 1500  
 caaccatggt gttggaattg tccacacaaa ctacttagag tacatcaaga gggagaagaa 1560  
 tggagctctt caagcatttt ttgtgaacca tgtaaacaat tgggtcacac gagcgtattg 1620  
 tgacaagggt cttcgctct ctgcggcaac acaagattta ccaaagtctg ttgtatgcaa 1680  
 tgtccatggt gtcaatccca agttccttat gattggggag aaaattgctg aagagagatc 1740  
 ccgtggtgaa caagctttct caaaagggtgc atacttctta ggaaaaatgg tgtgggctaa 1800  
 aggatacaga gaactaatag atctgatggc taaacacaaa agcgaacttg ggagcttcaa 1860  
 tctagatgta tatgggaacg gtgaagatgc agtcgaggtc caacgtgcag caaagaaaca 1920  
 tgacttgaat ctcaatttcc tcaaaggaag ggaccacgct gacgatgctc ttcacaagta 1980  
 caaagtgttc ataaacccca gcatcagcga tgttctatgc acagcaaccg cagaagcact 2040  
 agccatgggg aagtttgtgg tgtgtgcaga tcacccttca aacgaattct ttagatcatt 2100  
 cccgaactgc ttaacttaca aaacatccga agactttgtg tccaaagtgc aagaagcaat 2160  
 gacgaaagag ccactacctc tcaactcctga acaaatgtac aatctctctt gggaagcagc 2220  
 aacacagagg ttcattggagt attcagatct cgataagatc ttaaacaatg gagagggagg 2280  
 aaggaagatg cgaaaatcaa gatcgggtcc gagctttaa gaggtggtcg atggaggatt 2340  
 ggcattctca cactatgttc taacagggaa cgatttcttg agactatgca ctggagcaac 2400  
 accaagaaca aaagactatg ataatacaaa ttgcaaggat ctgaatctcg taccacctca 2460  
 cgttcacaaag ccaatcttcg gctggtagat atttcccat aggccacca gttattgctt 2520  
 gtgacttatt aaaccactac gttattgtta tcctttttta cttttacagt tgttgtaggt 2580  
 cgtttgtttg ttaatagaaa gggtagatta ttattaaaaa aaaaaaaaaa aaaaaaaaaa 2640  
 aaactcgag 2649

<210> 2  
 <211> 808  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 2  
 Met Val Lys Glu Thr Leu Ile Pro Pro Ser Ser Thr Ser Met Thr Thr  
 1 5 10 15  
 Gly Thr Ser Ser Ser Ser Ser Leu Ser Met Thr Leu Ser Ser Thr Asn  
 20 25 30  
 Ala Leu Ser Phe Leu Ser Lys Gly Trp Arg Glu Val Trp Asp Ser Ala  
 35 40 45  
 Asp Ala Asp Leu Gln Leu Met Arg Asp Arg Ala Asn Ser Val Lys Asn  
 50 55 60  
 Leu Ala Ser Thr Phe Asp Arg Glu Ile Glu Asn Phe Leu Asn Asn Ser  
 65 70 75 80  
 Ala Arg Ser Ala Phe Pro Val Gly Ser Pro Ser Ala Ser Ser Phe Ser  
 85 90 95

Asn	Glu	Ile	Gly	Ile	Met	Lys	Lys	Leu	Gln	Pro	Lys	Ile	Ser	Glu	Phe	100	105	110
Arg	Arg	Val	Tyr	Ser	Ala	Pro	Glu	Ile	Ser	Arg	Lys	Val	Met	Glu	Arg	115	120	125
Trp	Gly	Pro	Ala	Arg	Ala	Lys	Leu	Gly	Met	Asp	Leu	Ser	Ala	Ile	Lys	130	135	140
Lys	Ala	Ile	Val	Ser	Glu	Met	Glu	Leu	Asp	Glu	Arg	Gln	Gly	Val	Leu	145	150	155
Glu	Met	Ser	Arg	Leu	Arg	Arg	Arg	Arg	Asn	Ser	Asp	Arg	Val	Arg	Phe	165	170	175
Thr	Glu	Phe	Phe	Ala	Glu	Ala	Glu	Arg	Asp	Gly	Glu	Ala	Tyr	Phe	Gly	180	185	190
Asp	Trp	Glu	Pro	Ile	Arg	Ser	Leu	Lys	Ser	Arg	Phe	Lys	Glu	Phe	Glu	195	200	205
Lys	Arg	Ser	Ser	Leu	Glu	Ile	Leu	Ser	Gly	Phe	Lys	Asn	Ser	Glu	Phe	210	215	220
Val	Glu	Lys	Leu	Lys	Thr	Ser	Phe	Lys	Ser	Ile	Tyr	Lys	Glu	Thr	Asp	225	230	235
Glu	Ala	Lys	Asp	Val	Pro	Pro	Leu	Asp	Val	Pro	Glu	Leu	Leu	Ala	Cys	245	250	255
Leu	Val	Arg	Gln	Ser	Glu	Pro	Phe	Leu	Asp	Gln	Ile	Gly	Val	Arg	Lys	260	265	270
Asp	Thr	Cys	Asp	Arg	Ile	Val	Glu	Ser	Leu	Cys	Lys	Cys	Lys	Ser	Gln	275	280	285
Gln	Leu	Trp	Arg	Leu	Pro	Ser	Ala	Gln	Ala	Ser	Asp	Leu	Ile	Glu	Asn	290	295	300
Asp	Asn	His	Gly	Val	Asp	Leu	Asp	Met	Arg	Ile	Ala	Ser	Val	Leu	Gln	305	310	315
Ser	Thr	Gly	His	His	Tyr	Asp	Gly	Gly	Phe	Trp	Thr	Asp	Phe	Val	Lys	325	330	335
Pro	Glu	Thr	Pro	Glu	Asn	Lys	Arg	His	Val	Ala	Ile	Val	Thr	Thr	Ala	340	345	350
Ser	Leu	Pro	Trp	Met	Thr	Gly	Thr	Ala	Val	Asn	Pro	Leu	Phe	Arg	Ala	355	360	365
Ala	Tyr	Leu	Ala	Lys	Ala	Ala	Lys	Gln	Ser	Val	Thr	Leu	Val	Val	Pro	370	375	380
Trp	Leu	Cys	Glu	Ser	Asp	Gln	Glu	Leu	Val	Tyr	Pro	Asn	Asn	Leu	Thr	385	390	395
Phe	Ser	Ser	Pro	Glu	Glu	Gln	Glu	Ser	Tyr	Ile	Arg	Lys	Trp	Leu	Glu	405	410	415
Glu	Arg	Ile	Gly	Phe	Lys	Ala	Asp	Phe	Lys	Ile	Ser	Phe	Tyr	Pro	Gly	420	425	430
Lys	Phe	Ser	Lys	Glu	Arg	Arg	Ser	Ile	Phe	Pro	Ala	Gly	Asp	Thr	Ser	435	440	445
Gln	Phe	Ile	Ser	Ser	Lys	Asp	Ala	Asp	Ile	Ala	Ile	Leu	Glu	Glu	Pro	450	455	460

Glu His Leu Asn Trp Tyr Tyr His Gly Lys Arg Trp Thr Asp Lys Phe  
 465 470 475 480  
 Asn His Val Val Gly Ile Val His Thr Asn Tyr Leu Glu Tyr Ile Lys  
 485 490 495  
 Arg Glu Lys Asn Gly Ala Leu Gln Ala Phe Phe Val Asn His Val Asn  
 500 505 510  
 Asn Trp Val Thr Arg Ala Tyr Cys Asp Lys Val Leu Arg Leu Ser Ala  
 515 520 525  
 Ala Thr Gln Asp Leu Pro Lys Ser Val Val Cys Asn Val His Gly Val  
 530 535 540  
 Asn Pro Lys Phe Leu Met Ile Gly Glu Lys Ile Ala Glu Glu Arg Ser  
 545 550 555 560  
 Arg Gly Glu Gln Ala Phe Ser Lys Gly Ala Tyr Phe Leu Gly Lys Met  
 565 570 575  
 Val Trp Ala Lys Gly Tyr Arg Glu Leu Ile Asp Leu Met Ala Lys His  
 580 585 590  
 Lys Ser Glu Leu Gly Ser Phe Asn Leu Asp Val Tyr Gly Asn Gly Glu  
 595 600 605  
 Asp Ala Val Glu Val Gln Arg Ala Ala Lys Lys His Asp Leu Asn Leu  
 610 615 620  
 Asn Phe Leu Lys Gly Arg Asp His Ala Asp Asp Ala Leu His Lys Tyr  
 625 630 635 640  
 Lys Val Phe Ile Asn Pro Ser Ile Ser Asp Val Leu Cys Thr Ala Thr  
 645 650 655  
 Ala Glu Ala Leu Ala Met Gly Lys Phe Val Val Cys Ala Asp His Pro  
 660 665 670  
 Ser Asn Glu Phe Phe Arg Ser Phe Pro Asn Cys Leu Thr Tyr Lys Thr  
 675 680 685  
 Ser Glu Asp Phe Val Ser Lys Val Gln Glu Ala Met Thr Lys Glu Pro  
 690 695 700  
 Leu Pro Leu Thr Pro Glu Gln Met Tyr Asn Leu Ser Trp Glu Ala Ala  
 705 710 715 720  
 Thr Gln Arg Phe Met Glu Tyr Ser Asp Leu Asp Lys Ile Leu Asn Asn  
 725 730 735  
 Gly Glu Gly Gly Arg Lys Met Arg Lys Ser Arg Ser Val Pro Ser Phe  
 740 745 750  
 Asn Glu Val Val Asp Gly Gly Leu Ala Phe Ser His Tyr Val Leu Thr  
 755 760 765  
 Gly Asn Asp Phe Leu Arg Leu Cys Thr Gly Ala Thr Pro Arg Thr Lys  
 770 775 780  
 Asp Tyr Asp Asn Gln His Cys Lys Asp Leu Asn Leu Val Pro Pro His  
 785 790 795 800  
 Val His Lys Pro Ile Phe Gly Trp  
 805

<210> 3  
 <211> 2500

<212> DNA  
<213> *Arabidopsis thaliana*

<220>  
<221> misc\_feature  
<222> (467)..(983)  
<223> exon 1

<220>  
<221> misc\_feature  
<222> (1201)..(1427)  
<223> exon 2

<220>  
<221> misc\_feature  
<222> (1608)..(1874)  
<223> exon 3

<220>  
<221> misc\_feature  
<222> (1987)..(2073)  
<223> exon 4

<220>  
<221> misc\_feature  
<222> (2261)..(2332)  
<223> exon 5

<400> 3  
tgtgggatta agattctata ttatcttttt gataataaaa acaaaaatat tagcaaaata 60  
gaaaagattc acaagatctt tattagaaga aaagtgaatg gtatgcataa ataaaatcca 120  
tgaaaagtaa accattaatg cacaggacac gagttactca tagttcatat gttaaattgg 180  
ttttctatta gcaattagat atggctctaa tatgagcaat tggccacata agtggacaaa 240  
tgccaattcg aaatgatatg tacagcttct gattctttta atggaatcat atactaaatt 300  
ttataattca gactcgtgac atcatatcatg ttccacaaga tgaaaaatct aactttcatc 360  
atcaatatta caccaaagct gctatagaat gattcagata gtatacattg atttatacac 420  
aaaaatatcc caatgatatg cagacaaaga taaattaaaa aacagggtcaa tcttgcttgc 480  
gagtatttgg ggaaggagtg ttgagagaca aaccaagatc tctacataac tcttcacg 540  
gctgcaagct tccaggtata gcaccaaag ctgttctgga agcttcaa atccagacgcca 600  
agaaatgaat atacgctgac atatcctcca agtttttccc cacactaatc gaagatgacg 660  
caaacacact tctctttgac aagttcgaat cagctcttga taaccgggtc agatcagaga 720  
cttttataaa ccgttggtga gcagcttccc atgatagttc atgcctttgt tgctctgtaa 780  
gttgcgatgg ttgttcccca agagccttga gcgtggctct taaaaacct tgtccatcgt 840  
cataggttct gcagttggga aactgtttga agaacttggt cgatatgtga ttcgcgcata 900  
ctactatttt tcccatcgcc aaggcttctg cagttgttgt acacacaacg tctgtcgtgc 960  
tggggttgag aaacacttta tagctgaaaa tagaatggg aaaagttaa ggtatagaat 1020  
ccgatagact gcaaggaaat caatcaatct tgaggctgtt aagaactaga gacttacttg 1080  
tgaaatagcg agtcagcgtg atcacgtcct gggtaaacaat taaccgtcaa gtcgagtttt 1140  
cgggctgctt ctttgatctc ttcagagtc tctccatcac cgtataaatc aacctctaac 1200

tcggcaagtt ccttttggtg tttctcaagt agtttaagaa gtccttgta ccctttgctc 1260  
cagaccatct taccaatgta gtatgcgcct ttagtgaagg gctgctcctg gagcttctgc 1320  
tgttctagtt ttctcaaccc aatttcgaga aatttagggg tgacgccatg aacattgcag 1380  
actatagatt taggggtattc ttgagtcgca gcagataacc tgattacctg caggagtaac 1440  
atcattggta caagtcctga ttcacatcaa aaggaccggc cataatttg gtcacattcc 1500  
aacaaactgt cttgattctc aaattaaacc cagccattgg atatattcac ataattgagc 1560  
aatcactcta tcatgtccta gtggctgtaa agaagaaaga agtttacctt gtggcagtaa 1620  
atgccaaaca cccaactatt taagtatttg aggaaaaatg ctttgacacg gccttgtttc 1680  
tctcttttaa cgtattccaa gtagttagtg tgtacgattc ctatgacgta gttgaactta 1740  
gttttccatt tttggccatg atgaaacat gtgagatgct caggctcctc gaggacagca 1800  
atgtctgcct cttcatcagg aatggcatca gatatatccc caacaggaag aatactcctt 1860  
ttgtcaatag caaactgcc aagataaat ttgaattgct aagggtggca caacacagtc 1920  
acagagtact acgtaaaaga aatggaagaa ttcaactcta tatcacacgg aaatgaaaca 1980  
atcgactcta tgaaaggaag cagtatgtgt ttggtacca aaagaagatt cttgagagtc 2040  
atacctttcc aggatagaaa cgtatctcaa aggctaaacg aaaagagact ctctcctcaa 2100  
gccactggcg gacataagct tcttgctctg acgggggaact aaaagtgatg ctatttgggt 2160  
agacaagctt ttggtgcttc aaagtcagcc atggaatcac caacgtgacc cgtctttccc 2220  
catcatttgc aaggtaggca gcacggaaga gaggattaac agcagttccc gtaagccatg 2280  
gaatactage tgttgtaa atcgcaatgt gttgctcctg ctgattagtc atatcatctc 2340  
aagctaaacc tctacaaatt atcaaacaaa aagagtaaac caatagattc ttgtaattga 2400  
gtttgatctg attattgcat cttcccaata aggataacat tgcactacaa attcttaatt 2460  
tttctgcaaa ttcaaacaat cttttcacac gattcaagcg 2500

<210> 4

<211> 389

<212> PRT

<213> *Arabidopsis thaliana*

<400> 4

Met	Thr	Asn	Gln	Gln	Glu	Gln	His	Ile	Ala	Ile	Phe	Thr	Thr	Ala	Ser
1				5					10					15	

Ile	Pro	Trp	Leu	Thr	Gly	Thr	Ala	Pro	Leu	Arg	Tyr	Val	Ser	Ile	Leu
		20						25					30		

Glu	Arg	Tyr	Asp	Ser	Gln	Glu	Ser	Ser	Phe	Gly	Tyr	Gln	Thr	His	Thr
	35						40					45			

Ala	Ser	Phe	His	Arg	Phe	Ala	Ile	Asp	Lys	Arg	Ser	Ile	Leu	Pro	Val
	50					55					60				

Gly	Asp	Ile	Ser	Asp	Ala	Ile	Pro	Asp	Glu	Glu	Ala	Asp	Ile	Ala	Val
65					70					75					80

Leu	Glu	Glu	Pro	Glu	His	Leu	Thr	Trp	Phe	His	His	Gly	Gln	Lys	Trp
				85					90					95	

Lys Thr Lys Phe Asn Tyr Val Ile Gly Ile Val His Thr Asn Tyr Leu  
 100 105 110  
 Glu Tyr Val Lys Arg Glu Lys Gln Gly Arg Val Lys Ala Phe Phe Leu  
 115 120 125  
 Lys Tyr Leu Asn Ser Trp Val Val Gly Ile Tyr Cys His Lys Val Ile  
 130 135 140  
 Arg Leu Ser Ala Ala Thr Gln Glu Tyr Pro Lys Ser Ile Val Cys Asn  
 145 150 155 160  
 Val His Gly Val Asn Pro Lys Phe Leu Glu Ile Gly Leu Arg Lys Leu  
 165 170 175  
 Glu Gln Gln Lys Leu Gln Glu Gln Pro Phe Thr Lys Gly Ala Tyr Tyr  
 180 185 190  
 Ile Gly Lys Met Val Trp Ser Lys Gly Tyr Lys Glu Leu Leu Lys Leu  
 195 200 205  
 Leu Glu Lys His Gln Lys Glu Leu Ala Asp Tyr Lys Val Phe Leu Asn  
 210 215 220  
 Pro Ser Thr Thr Asp Val Val Cys Thr Thr Thr Ala Glu Ala Leu Ala  
 225 230 235 240  
 Met Gly Lys Ile Val Val Cys Ala Asn His Ile Ser Asn Lys Phe Phe  
 245 250 255  
 Lys Gln Phe Pro Asn Cys Arg Thr Tyr Asp Asp Gly Gln Gly Phe Val  
 260 265 270  
 Arg Ala Thr Leu Lys Ala Leu Gly Glu Gln Pro Ser Gln Leu Thr Glu  
 275 280 285  
 Gln Gln Arg His Glu Leu Ser Trp Glu Ala Ala Thr Gln Arg Phe Ile  
 290 295 300  
 Lys Val Ser Asp Leu Asn Arg Leu Ser Arg Ala Asp Ser Asn Leu Ser  
 305 310 315 320  
 Lys Arg Ser Val Phe Ala Ser Ser Ser Ile Ser Val Gly Lys Asn Leu  
 325 330 335  
 Glu Asp Met Ser Ala Tyr Ile His Phe Leu Ala Ser Gly Phe Glu Ala  
 340 345 350  
 Ser Arg Thr Ala Phe Gly Ala Ile Pro Gly Ser Leu Gln Pro Asp Glu  
 355 360 365  
 Glu Leu Cys Arg Asp Leu Gly Leu Ser Leu Asn Thr Pro Ser Pro Asn  
 370 375 380  
 Thr Arg Lys Gln Asp  
 385

<210> 5  
 <211> 5128  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> misc feature  
 <222> (627)..(1369)  
 <223> exon 1

<220>

<221> misc feature  
<222> (1565)..(1597)  
<223> exon 2

<220>  
<221> misc feature  
<222> (1712)..(1798)  
<223> exon 3

<220>  
<221> misc feature  
<222> (2186)..(2668)  
<223> exon 4

<220>  
<221> misc feature  
<222> (2769)..(3035)  
<223> exon 5

<220>  
<221> misc feature  
<222> (3137)..(3487)  
<223> exon 6

<220>  
<221> misc feature  
<222> (3570)..(4207)  
<223> exon 7

<220>  
<221> gene  
<222> (674)..(4079)  
<223> coding region for DGD1

<400> 5  
aagctttaac gagagcgatc aaaggcaaact cactagaaac tcttgagcaa gagttcatgt 60  
cgaccgtgat attttgtatg atatatatat ggtcaagatg gtgaagacaa aggttctact 120  
attgtataat ggtatcaaaa ttgtctaaat ttctaacaat ttcacaaggc ctgcgctggc 180  
gtccggttggc tgttttgggt aagtcggcct ctattttgat ccgttaatgg aactcgttac 240  
ggtctaatta cgtccactga taccactttc caatgttttt ttttgttaaa tgtcattgtc 300  
aataccatgc ccgtaggac cagattcttc gattccagca tattcctaag ttaaattccga 360  
cacaatcgga taatctcgtc aaccactcac aacacgccac gtaatcaatg atgtggaacc 420  
catgccacgt tggacacggc ctacacgggac aaggcttcag taaagagcgc ctatcgtcgc 480  
tccgacttgt ttattttcca cgtgttaatc ttccagaata atcagaaaga gaattaaaaa 540  
aataaaaacaa tccaaattaa tctcagccgt caattttctt tcttcttctt cccaaattct 600  
ctcaacagat agagaaaacc ttattagcag cactctcacg aaatcgtcgt gacgaaacga 660  
taaacacctaa gtcattgtaa aggaaactct aattcctccg tcatctacgt caatgacgac 720  
cggaacatct tcttcttcgt ctctttcaat gacgttatcc tcaacaaacg cgttatcggt 780  
tttgtcgaac ggatggagag aggtatggga ttcagcagat gcggatttgc agctgatgcg 840  
agacagagct aactctgtta agaattctagc atcaacgttc gatagagaga tcgagaattt 900  
cctcaataac tcggcgaggc ctgcgtttcc cgttggttca ccatcggcgt cgtctttctc 960  
aatgaaatt ggtatcatga agaagcttca gccgaagatt tcggagtctc gtaggggtta 1020

ttcggcgccg gagattagtc gcaaggttat ggagagatgg ggacctgcga gagcgaagct 1080  
tggaatggat ctatcggcga ttaagaaggc gatttgtctt gagatggaat tggatgagcg 1140  
tcagggagtt ttggagatga gtagattgag gagacggcgt aatagtataa gggtttaggtt 1200  
tacggagttt ttcgaggagg ctgagagaga tggagaagct tatttcggtg attgggaacc 1260  
gattaggtct ttgaagagta gatttaaaga gtttgagaaa cgaagctcgt tagaaatatt 1320  
gagtggattc aagaacagtg aatttggtga gaagctcaaa accagctttg taagtttctc 1380  
caactttttg ggaacctatt tcaaaagttt cttctatctt actgtagaag tggcttctct 1440  
ttcaatagcc acgacatttt cgtatgctga cttgatagtt actgttcttg gaatgattga 1500  
ttgagtttct tatgggtgtt ggcaactttt ctaatggtgc ttttttcttt tgttttattt 1560  
gcagaaatca atttacaaag aaactgatga ggctaagggt ggtatttgga agttggagct 1620  
ctatcagttt ttcttggtac tttatatatt ttaggtcga gttcagtggt ggattgggag 1680  
tttaacttgg cctcctttta ctttggtgca ggatgtccct ccgttggtg tacctgaact 1740  
gttggcatgt ttggttagac aatctgaacc tttcttgat cagattggtg ttagaaaggg 1800  
taagattgca tttttctctt catgatggtt aattattttg tctgttggtt gtattgagtt 1860  
gtttctctac cataggtggt ttttcggtca aaagtttgaa tcttctctct gatattagag 1920  
tgtctttgtt aagtgggttg cttgcttcac cagaagttta gatggtgaga tttgatgttc 1980  
tgcattatca cactaatcct ggatatcaaa tatgtgtaag tctagattct gtatgagaca 2040  
gatcaatcaa atgacatctg ccgtagacat aaaaatttct agatgtgtag gttattgggt 2100  
ttaaacaccc tttcttgtag attatcttat agtttcagtg tttacatata aagttcctga 2160  
ttctgttctg ctgaattttc tttcagatac atgtgaccga atagtagaaa gcctttgcaa 2220  
atgcaagagc caacaacttt ggcgctctgc atctgcacaa gcatccgatt taattgaaaa 2280  
tgataaccat ggagttgatt tggatatgag gatagccagt gttcttcaaa gcacaggaca 2340  
ccattatgat ggtgggtttt ggactgattt tgtgaagcct gagacaccgg aaaacaaaag 2400  
gcatgtggca attgttacia cagctagtct tcttggtg agcgggaacag ctgtaaatcc 2460  
gctattcaga gcggcgtatt tggcaaaagc tgcaaaacag agtgttactc tcgtgggttc 2520  
ttggctctgc gaatctgac aagaactagt gtatccaaac aatctcacct tcagctcacc 2580  
tgaagaacaa gagagttata tacgtaaagtg gttggaggaa aggattgggt tcaaggctga 2640  
ttttaaaatc tccttttacc caggaaaggt atgttgatca ttttggttct tattttttta 2700  
tttctatggc tgccaatatg tttttcaatt atttctatag agtaactgag ctttctgggt 2760  
tcttatagtt ttcaaaagaa aggcgcagca tatttcctgc tggtagacct tctcaattta 2820  
tatcgtcaaa agatgctgac attgctatac ttgaagaacc tgaacatctc aactgggtatt 2880  
atcacggcaa gcgttggtgact gataaattca accatgttgt tgggaattgtc cacacaaact 2940  
acttagagta catcaagagg gagaagaatg gagctcttca agcatttttt gtgaaccatg 3000  
taaacaattg ggtcacacga gcgtattgtg acaagggtga tcatctactc tatttcttca 3060  
agccttggtc tgttgcttga atcctcttta ctaataaata gtacacgagc taatacatat 3120

tttctactca tgaaagggttc ttgcctcttc tgcggcaaca caagatttac caaagtctgt 3180  
tgtatgcaat gtccatggtg tcaatcccaa gttccttatg attggggaga aaattgctga 3240  
agagagatcc cgtggtgaac aagctttctc aaaagggtgca tacttcttag gaaaaatggg 3300  
gtgggctaaa ggatacagag aactaataga tctgatggct aaacacaaaa gcgaacttgg 3360  
gagcttcaat ctagatgtat atgggaacgg tgaagatgca gtcgaggtcc aacgtgcagc 3420  
aaagaaacat gacttgaatc tcaatttcct caaaggaagg gaccacgctg acgatgctct 3480  
tcacaagtaa gttctgaaaa atgtgctttg cttttaaaaa cttgttaagg ttctgctctt 3540  
tgattgtctt tcccacatct tgatgaagg acaaagtgtt cataaacccc agcatcagcg 3600  
atgttctatg cacagcaacc gcagaagcac tagccatggg gaagtttggt gtgtgtgcag 3660  
atcacccttc aaacgaattc tttagatcat tcccgaactg cttacttac aaaacatccg 3720  
aagactttgt gtccaaagtg caagaagcaa tgacgaaaga gccactacct ctactcctg 3780  
aacaaatgta caatctctct tgggaagcag caacacagag gttcatggag tattcagatc 3840  
tcgataagat cttaaacaat ggagagggag gaaggaagat gcgaaaatca agatcgggtc 3900  
cgagctttaa cgagggtggt gatggaggat tggcattctc acactatggt ctaacaggga 3960  
acgatttctt gagactatgc actggagcaa caccaagaac aaaagactat gataatcaac 4020  
attgcaagga tctgaatctc gtaccacctc acgttcacaa gccaatcttc ggctggtaga 4080  
tatttcccca taggccacct agttattgct tgtgacttat taaaccacta cgttattggt 4140  
atcctttttt acttttacag ttgttgtagg tcgtttgttt gttaatagaa agggtagatt 4200  
attattagat gtctttttgt aaaatatcaa tacgaagcgt atttgatgat atataaaata 4260  
actatattgg caaaaatatg aactatgaag gccgttttcg tgattttggt cttttgtttc 4320  
acgaattcaa gctattcccc tttttttacg ccaaagatga aaagaacctt ccctattaat 4380  
atcgctattg tctaaaattt cgaaaactac tttaatcacg actagaccaa atatatgtcg 4440  
accgataccg atagagaaat tagtgccccg tctaatactt tctctccaaa attacagaat 4500  
atttagagta gttaatcaac gtaacacgac aaggaaaatg atggaaaaag tgggtggtttc 4560  
tgctttggca actagtgtta ggtcacttac gtctctttt tctgtattgg aaattacgtg 4620  
gataaattga actttctttc aatctctatc aaattattaa tccacacatg tatacgcaat 4680  
atatgatcat taaataaata aaaaggtaga ttggtctata aattcgtatc acaaattggac 4740  
taataatttg tagtgaaaac tcatttaccc atgtgacagc tccaaatttc tgaacttttt 4800  
attttgaggg atggtacaaa tccgagttcc atgatcatgg aaaaatcaaa tttacaaaac 4860  
acaaattact gtttgaaaca agcaagttac tatatatgta gtttgatttc acactagaga 4920  
atctactgat aatgaatttt ttatatatcg tgaagctgaa agtgaaatta taactagcta 4980  
gttgaattgc ttattatggg tggaggggag ccgatgaaaa attcttcgac cacacataaa 5040  
gtcacccttc taaagaacac ttgcaccgac cacatcaatc acgcattcat aaattttcaa 5100  
catttatata aatgtaggaa aaaaacaa 5128

<210> 6  
<211> 5050  
<212> DNA  
<213> *Arabidopsis thaliana*

<220>  
<221> mutation  
<222> (334)  
<223> T to C exchange (5' non-transcribed region)

<220>  
<221> mutation  
<222> (3189)  
<223> C to T exchange (CAA to TAA, premature stop codon)

<220>  
<221> mutation  
<222> (3434)  
<223> G insertion after 3434 (silent, in intron)

<220>  
<221> mutation  
<222> (3649)  
<223> C to T exchange (conservative exchange, Ser codon)

<220>  
<221> mutation  
<222> (4795)  
<223> A insertion after 4795 (3' non-transcribed region)

<220>  
<221> mutation  
<222> (4868)  
<223> T to A exchange (3' non-transcribed region)

<220>  
<221> mutation  
<222> (4874)  
<223> A to T exchange (3' non-transcribed region)

<400> 6

```
ttttgtatga tatatatatg gtcaagatgg tgaagacaaa ggttctacta ttgtataatg 60
gtatcaaaat ttgctaaatt tctaacaatt tcacaagggtc tcgctgggcg tccgttggtt 120
gttttggtta agtcggcctc tattttgatc cgттаатgga actcggttacg gtctaattac 180
gtccactgat atcactttcc aatgtttttt tttgttaaаt gtcattgtca ataccatgcc 240
ccgtaggacc agattcttcg attccagcat attcctaagt taaatccgac acaatcggat 300
aatctcgtca accactcaca acacgccacg taaccaatga tgtggaaccc atgccacgtt 360
ggacacgggtc tcacgggaca aggcttcagt aaagagcgcc tatcggtccgt ccgacttggt 420
tattttccac gtgttaatct tccagaataa tcagaaagag aattaaaaaa ataaaacaat 480
ccaaattaat ctсagccgtc aattttctctt cttctctcttc ccaaattctc tcaacagata 540
gagaaaacct tattagcagc actctcacga aatcgtcgtg acgaaacgat aaaccctaag 600
tcatggtaaa ggaaactcta attcctccgt catctacgtc aatgacgacc ggaacatctt 660
cttcttcgtc tctttcaatg acgttatcct caacaaacgc gttatcgttt ttgtcgaaag 720
gatggagaga ggtatgggat tcagcagatg cggatttgca gctgatgсga gacagagcta 780
```

actctgttaa gaatctagca tcaacgttcg atagagagat cgagaatttc ctcaataact 840  
cggcgaggtc tgcgtttccc gttgggtcac catcggcgtc gtctttctca aatgaaattg 900  
gtatcatgaa gaagcttcag ccgaagattt cggagtttcg tagggtttat tggcgccgg 960  
agattagtcg caaggttatg gagagatggg gacctgagag agcgaagctt ggaatggatc 1020  
tateggcgat taagaaggcg attgtgtctg agatggaatt ggatgagcgt caggaggtt 1080  
tggagatgag tagattgagg agacggcgta atagtatag ggtaggtt acggagttt 1140  
tcgcggaggc tgagagagat ggagaagctt atttcggtga ttgggaaccg attaggctt 1200  
tgaagagtag atttaaagag ttgagaaac gaagctcgtt agaaatattg agtggattca 1260  
agaacagtga atttggtgag aagctcaaaa ccagctttgt aagtttctcc aactttttgg 1320  
gaacctatth caaaagtttc ttctatctta ctgtagaagt ggcttctctt tcaatagcca 1380  
cgacatthtc gtatgctgac ttgatagtta ctgttcttgg aatgattgat tgagtttctt 1440  
atggtgtttg gcaactthtc taatggtgct ttttctttt gttttatttg cagaaatcaa 1500  
tttacaaga aactgatgag gctaagggtg gtatttgaa gttggagctc tatcagttt 1560  
tcttgttact ttatatattt ttaggtcgag ttcagtgtt gattgggagt ttaacttggc 1620  
ctccttttac tttggtgcag gatgtccctc cggtggatgt acctgaactg ttggcatgtt 1680  
tggttagaca atctgaacct tttcttgatc agattgggtg tagaaagggt aagattgcat 1740  
tttctcttc atgatggtta attattttgt ctgttgatg tattgagttg tttctctacc 1800  
atagggtggt tttccgtcaa aagtttgaat cttctctctg atattagagt gtctttgtta 1860  
agtgggttgc ttgcttcacc agaagtttag atggtgagat ttgatgttct gcattatcac 1920  
actaatcctg gatatcaaat atgtgtaagt ctagattctg tatgagacag atcaatcaaa 1980  
tgacatctgc cgtagacata aaaatttcta gatgtgtagg ttattgggtt taaacaccct 2040  
ttcttgtaaa ttatcttata gtttcagtgt ttacatacaa agttcctgat tctgttctgc 2100  
tgaattttct ttcagataca tgtgaccgaa tagtagaaag cttttgcaa tgcaagagcc 2160  
aacaactttg gcgtctgcca tctgcacaag catccgattt aattgaaaat gataaccatg 2220  
gagttgattt ggatatgagg atagccagtg ttttcaaag cacaggacac cattatgatg 2280  
gtgggttttg gactgatttt gtgaagcctg agacaccgga aaacaaaagg catgtggcaa 2340  
ttgttacaac agctagtctt ccttggatga ccggaacagc tgtaaatecg ctattcagag 2400  
cggcgtatth ggcaaaagct gcaaaacaga gtgttactct cgtgggttct tggtctctgc 2460  
aatctgatca agaactagt tatccaaaca atctcacct cagctcacct gaagaacaag 2520  
agagttatat acgtaaatgg ttggaggaaa ggattgggtt caaggctgat tttaaaatct 2580  
ccttttacc aggaaaggta tgttgatcat tttggattct atttttttat ttctatggct 2640  
gccaatatgt ttttcaatta tttctataga gtaactgagc tttctgggtt cttatagttt 2700  
tcaaaagaaa ggcgcagcat atttctgtct ggtgacact ctcaatttat atcgtcaaaa 2760  
gatgctgaca ttgctatact tgaagaacct gaacatctca actggtatta tcacggcaag 2820

cgttggactg ataaattcaa ccatgttggt ggaattgtcc acacaaacta cttagagtac 2880  
atcaagaggg agaagaatgg agctcttcaa gcattttttg tgaaccatgt aaacaattgg 2940  
gtcacacgag cgtattgtga caaggtgaat catctactct atttcttcaa gccttggtct 3000  
gttgcttgaa tcctctttac taataaatag tacacgagct aatacatatt ttctactcat 3060  
gaaaggttct tcgcctctct gcggcaacac aagatttacc aaagtctgtt gtatgcaatg 3120  
tccatgggtg caatcccaag ttccttatga ttggggagaa aattgctgaa gagagatccc 3180  
gtgggtgaata agctttctca aaaggtgcat acttcttagg aaaaatgggtg tgggctaaag 3240  
gatacagaga actaatagat ctgatggcta aacacaaaag cgaacttggg agcttcaatc 3300  
tagatgtata tgggaacggg gaagatgcag tcgaggtcca acgtgcagca aagaaacatg 3360  
acttgaatct caatttcctc aaaggaaggg accacgctga cgatgctctt cacaagtaag 3420  
ttctgaaaaa tgtggctttg cttttaaaaa cttgttaagg tttcgtctct tgattgtctt 3480  
tcccacatct tgatgaaggt acaaagtgtt cataaacccc agcatcagcg atgttctatg 3540  
cacagcaacc gcagaagcac tagccatggg gaagtttgtg gtgtgtgcag atcacccttc 3600  
aaacgaattc tttagatcat tcccgaactg cttaacttac aaaacatctg aagactttgt 3660  
gtccaaagtg caagaagcaa tgacgaaaga gccactacct ctcaactctg aacaaatgta 3720  
caatctctct tgggaagcag caacacagag gtcatggag tattcagatc tcgataagat 3780  
cttaaacaat ggagagggag gaaggaagat gcgaaaatca agatcggttc cgagctttaa 3840  
cgaggtgggt gatggaggat tggcattctc acactatgtt ctaacagggg acgatttctt 3900  
gagactatgc actggagcaa caccaagaac aaaagactat gataatcaac attgcaagga 3960  
tctgaatctc gtaccacctc acgttcacaa gccaatcttc ggctggtaga tatttcccca 4020  
taggccaccc agttattgct tgtgacttat taaaccacta cgttattgtt atcctttttt 4080  
acttttacag ttgttgtagg tcgtttgttt gttaatagaa agggtagatt attattagat 4140  
gtctttttgt aaaatatcaa tacgaagcgt atttgatgat atataaata actatattgg 4200  
caaaaatatg aactatgaag gccgttttcg tgattttgtt cttttgtttc acgaattcaa 4260  
gctattcccc tttttttacg ccaaagatga aaagaacctt ccctattaat atcgctattg 4320  
tctaaaattt cgaaaactac ttaatcacg actagaccaa atatatgtcg accgataccg 4380  
atagagaaat tagtgccccg tctaatactt tctctccaaa attacagaat atttagagta 4440  
gttaatcaac gtaacacgac aaggaaaatg atggaaaaag tgggtggttc tgctttggca 4500  
actagtgtta ggtcacttac gtctctttt tctgtattgg aaattacgtg gataaattga 4560  
actttctttc aatctctatc aaattattaa tccacacatg tatacgcaat atatgatcat 4620  
taaataaata aaaagttaga ttgggtctata aattcgtatc acaaattggac taataatttg 4680  
tagtgaaaac tcattttacc atgtgacagc tccaaatttc tgaacttttt attttgaggg 4740  
atggtacaaa tccgagttcc atgatcatgg aaaaatcaaa ttttaacaaac acaaaattac 4800  
tgtttgaaac aggcaagtta ctatatatgt agtttgattt cacactagag aatctactga 4860  
taatgaaatt tttttatata gtgaagctga aagtgaattt ataactagct agttgaattg 4920

cttattatgg ttggagggga gccgatgaaa aattcttcga ccacacataa agtcaccttt 4980  
ctaaagaaca cttgcaccga ccacatcaat cagcattca taaattttca acatttatat 5040  
aaatgtagga 5050

<210> 7  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Oligonucleotide  
primer

<400> 7  
gcggatccgg taaaggaaac tctaatt 27

<210> 8  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Oligonucleotide  
primer

<400> 8  
ttctgcagtc taccagccga agattgg 27

**WE CLAIM:**

1. An isolated nucleic acid molecule comprising a nucleic acid sequence encoding for a polypeptide having digalactosyldiacylglycerol galactosyltransferase activity.
2. The nucleic acid molecule of Claim 1 wherein the nucleic acid molecule has a nucleotide sequence comprising SEQ ID NO: 1.
3. The isolated nucleic acid molecule of Claim 1 wherein the nucleotide sequence encodes a polypeptide having an amino acid sequence comprising SEQ ID NO: 2.
4. A vector comprising the nucleic acid of Claim 1.
5. A host cell comprising the vector of Claim 4.
6. The host cell of Claim 5 wherein the host cell is a plant cell.
7. An isolated nucleic acid molecule comprising a nucleotide sequence capable of hybridizing under high stringency conditions to SEQ ID NO: 1 or the complement of SEQ ID NO: 1.
8. The nucleic acid molecule of Claim 7, wherein the nucleic acid molecule is capable of changing the lipid composition of a plant cell.
9. A vector comprising the nucleic acid of Claim 7.
10. A host cell comprising the vector of Claim 9.
11. The host cell of Claim 10 wherein the host cell is a plant cell.

12. A method for producing a polypeptide encoded by the nucleic acid molecule of Claim 1 comprising the steps of:

- a) inserting the nucleic acid molecule into a vector;
- b) transforming a host cell with the vector; and
- c) growing a culture of the host cell in a suitable culture medium.

13. The method of Claim 12 wherein the host cell is *E. coli*.

14. The method of Claim 12 wherein the host cell is yeast.

15. A polypeptide comprising digalactosyldiacylglycerol galactosyltransferase activity.

16. The polypeptide of Claim 15 wherein the polypeptide has an amino acid sequence comprising SEQ ID NO: 2.

17. The polypeptide of Claim 15 wherein the polypeptide is encoded for by a nucleic acid molecule having a nucleotide sequence comprising SEQ ID NO: 1.

18. The polypeptide of Claim 15 wherein the polypeptide is encoded for by a nucleic acid molecule having a nucleotide sequence comprising SEQ ID NO: 3.

19. A method for altering leaf lipid composition in a plant comprising introducing a nucleic acid molecule encoding for a polypeptide comprising digalactosyldiacylglycerol galactosyltransferase activity into a plant.

20. The method of Claim 19 wherein the nucleic acid molecule comprises SEQ ID NO: 1.

21. The method of Claim 19 wherein the nucleic acid molecule comprises SEQ ID NO: 3.

22. A transgenic plant produced by the method of Claim 19.

23. The seeds of the plant of Claim 22.
24. A transgenic plant produced by breeding the plant of Claim 22 wherein the plant retains the trait of altered leaf lipid composition.
25. A method for determining inhibitors of digalactosyldiacylglycerol galactosyltransferase comprising the steps of:
- a) contacting the galactosyltransferase with a compound; and
  - b) determining any change in activity of the galactosyltransferase.
26. The method of Claim 25 wherein the digalactosyldiacylglycerol galactosyltransferase has an amino acid sequence comprising SEQ ID NO: 2.
27. The method of Claim 25 wherein the digalactosyldiacylglycerol galactosyltransferase has an amino acid sequence comprising SEQ ID NO: 4.





Figure 1



Wild Type

*act1*

*dgd1*

*act1,dgd1*

Figure 2

Figure 3A




Figure 3B




Figure 3C






Figure 3D





**Figure 4**